

R. de la Torre and M. Dierssen (Eds.)

Progress in Brain Research, Vol. 197

ISSN: 0079-6123

Copyright © 2012 Elsevier B.V. All rights reserved.

CHAPTER 11

c0055

A Sonic hedgehog (Shh) response deficit in trisomic cells may be a common denominator for multiple features of Down syndrome

Duane G. Currier, Renita C. Polk and Roger H. Reeves*

*Department of Physiology, The McKusick-Nathans Institute for Genetic Medicine,
Johns Hopkins University School of Medicine, Baltimore, Maryland, USA*

Abstract: The hedgehog (HH) family of growth factors is involved in many aspects of growth and development, from the establishment of left–right axes at gastrulation to the patterning and formation of multiple structures in essentially every tissue, and the maintenance and regulation of stem cell populations in adults. Sonic hedgehog (Shh) in particular acts as a mitogen, regulating proliferation of target cells, a growth factor that triggers differentiation in target populations, and a morphogen causing cells to respond differently based on their positions along a spatial and temporal concentration gradient. Given its very broad range of effects in development, it is not surprising that many of the structures affected by a disruption in Shh signaling are also affected in Down syndrome (DS). However, recent studies have shown that trisomic cerebellar granule cell precursors have a deficit, compared to their euploid counterparts, in their response to the mitogenic effects of Shh. This deficit substantially contributes to the hypoplastic cerebellum in mouse models that parallels the human DS phenotype and can be corrected in early development by a single exposure to a small-molecule agonist of the Shh pathway.

Here, we consider how an attenuated Shh response might affect several aspects of development to produce multiple phenotypic outcomes observed in DS.

Keywords: SHH signaling; Down syndrome; brain development; common denominators of Down syndrome; neural crest; cerebellum.

Au3 s0005 Therapeutic approaches in Down syndrome

p0100 Trisomy for human chromosome 21 (Hsa21) results in Down syndrome (DS) which is among the most complex genetic perturbations compati-

ble with survival past term. While trisomy affects development of every tissue, reduced cognitive ability in DS is among the most limiting features, and DS is one of the leading genetic causes of intellectual disability. The development and characterization of mouse models of DS, especially Ts65Dn, demonstrates that orthologous gene dosage effects produce comparable outcomes for

Au1 *Corresponding author.
E-mail: rreeves@jhmi.edu

DOI: 10.1016/B978-0-444-54299-1.00011-X

195

some phenotypes, including cognitive impairment (Fernandez et al., 2007; Hanson et al., 2007; Kleschevnikov et al., 2004; Reeves et al., 1995). As detailed elsewhere in this volume (de la Torre and Dierssen, 2012; Fillat and Altafaj, 2012; Mobley, in this volume), several drugs with the potential to ameliorate cognitive deficits in DS are making their way to clinical trials.

[Au4]

p0105

Studies of mice have played an important role in understanding the brain regions that are especially affected in DS (Lott and Dierssen, 2010). Functional outcomes as well as anatomical and physiological studies demonstrate three regions among those with the largest effects: prefrontal cortex, a contributor to executive function; hippocampus, a crucial site for learning and memory; and cerebellum, which shows a dramatic reduction in size and cellularity. It is notable that initial studies of learning and memory deficits in trisomic mice, which showed affects on hippocampus, informed the development of the first cognitive tests focused on deficits associated with the hippocampus in DS (Pennington et al., 2003; Reeves et al., 1995). That effort has been carried forward, resulting recently in the Arizona Cognitive Test Battery for DS (ACTB) (Edgin et al., 2010). The ACTB is a sensitive set of tests focused on brain regions affected in DS (see Edgin et al., 2012). Clinical trials with the goal of ameliorating cognitive deficits in DS have begun; many proposed efforts with this goal will utilize the ACTB tests as part of their assessments (for updated information, see <http://clinicaltrials.gov/>).

[Au5]

p0110

Currently, approaches to therapy in DS may be thought of in three very broad areas. First, people with DS frequently exhibit early onset of geriatric diseases. The histopathology of Alzheimer disease (AD) is present in all persons with DS along with the sequellae of the disease, including dementia in a substantial fraction of the DS population, and is certainly related at least in part to overexpression of the amyloid

precursor protein gene, APP (Salehi et al., 2006). Age-related loss of afferents to the hippocampus from the locus coeruleus of neurons that use norepinephrine as a neurotransmitter and degeneration of basal forebrain cholinergic neurons are also hallmarks of DS shared with AD (Salehi et al., 2006, 2009).

A second general area for DS therapy involves correction of perturbed neuronal function in older children or adults. For example, restoration of an imbalance of inhibitory and excitatory inputs to the hippocampus forms the basis for major clinical trials going forward (Braudeau et al., 2011; Fernandez et al., 2007). This approach is based on the observation that downregulation of the GABAergic inhibitory PV neurons in Ts65Dn mice restores the balance of inhibitory:excitatory inputs and normalizes performance in hippocampal-based tasks such as the Novel Object Recognition Task and the Morris Water Maze (see Reeves and Garner, 2007; Rueda et al., 2008; Salehi et al., 2007). Several other efforts that have been carried out in trisomic mice and in some cases piloted in human studies look at a variety of hippocampal pathways (Lott and Dierssen, 2010).

p0115

A third potential area for therapy that is further downstream in the drug development pipeline addresses the initial basis of cognitive deficits, that is, antenatal brain development (Haydar and Reeves, 2012). Anatomical and morphological changes in the developing trisomic brain are being studied in detail in animal models, while imaging techniques are increasingly providing information about development of the DS brain. One approach of this type has been shown to normalize early deficits in postnatal development of the cerebellum, which is markedly hypocellular in DS and mouse models (Roper et al., 2006b); this example involves Shh signaling and is considered in detail here.

p0120

s0010 Shh signaling

s0015 Canonical Shh pathway


p0125 The Shh ligand is produced in cells distinct and often separated from those receiving the signal. The precursor protein is substantially modified by a cleavage that involves addition of a cholesterol moiety followed by palmitoylation (Mann and Beachy, 2004). Fully processed Shh (Shh-Np) is secreted from the producing cell and likely assembles into multimers (Zeng et al., 2001). Extracellular Shh-Np is sensed by the receiving cell via interactions with the 12-pass transmembrane protein, Patched (Ptch) (Marigo et al., 1996; Stone et al., 1996). In the pathway-off state (Fig. 1a), Ptch catalyzes the production of an unidentified repressor of Smoothened (Smo), a seven-pass transmembrane protein with possible G-protein-coupled receptor activity (Ayers and Therond, 2010; Chen et al., 2002). When Smo is repressed (pathway-off), the transcription factors Gli2 and Gli3 are targeted to the proteasome for processing to produce their transcriptional repressor forms (Gli2^R, Gli3^R) (Asai et al., 2006; Wang et al., 2000).

Another pathway element, Suppressor of Fused (SuFu), is found in both the cytoplasm and the nucleus and interacts with Gli1 and Gli2 proteins to further suppress pathway activity (Barnfield et al., 2005; Kogerman et al., 1999). SuFu/Gli complexes are exported from the nucleus and tethered in a SuFu-dependent manner in the cytoplasm. Further, SuFu inhibits Gli-mediated transcriptional activation by binding and inhibiting DNA-bound Gli1 or Gli2. The pathway is activated when Shh binds to Ptch, inhibiting the catalytic activity of the latter, thereby reversing the repression on Smo (Fig. 1b). This results in degradation of SuFu and Gli phosphorylation to produce activator Gli proteins that move to the nucleus and promote transcription (Chen et al., 2002; Humke et al., 2010; Yue et al., 2009; Zhang et al., 2004).

Noncanonical Shh signaling

s0020

Jenkins (2009) broadly defines several mechanisms p0135 for pathway activation outside of the canonical derepression of Gli transcription factors following Shh binding to Ptch. For example, Ptch can interact directly with CyclinB1 to affect cell cycle



f0005 Fig. 1. The Shh pathway. (a) In the pathway-off state, Patched (Ptch) catalytically inhibits Smoothened (Smo) activity through an unidentified intermediate. Suppressor of Fused (SuFu) mediates cleavage of Gli2 and Gli3 into their repressor forms, lacking transactivation domains. The Gli2/3 repressors translocate to the nucleus where they repress transcription by binding target gene promoter sequences. (b) Shh binding to Ptch inhibits its catalytic repression of Smo, resulting in activation of Smo and degradation of SuFu. In the absence of SuFu, Gli2/3 are phosphorylated to become the activator forms. Gli2/3 activators translocate to the nucleus where they promote transcription by recruiting other transcriptional activators to target gene promoters. (For color version of this figure, the reader is referred to the Web version of this chapter.)

progression (Barnes et al., 2001) and can initiate apoptosis in neuroepithelial cells until it is blocked by Shh binding (Thibert et al., 2003). Although Ptch is the primary receptor for Shh, several other membrane bound proteins compete for Shh and are capable of enhancing or inhibiting pathway activity. Cell Adhesion Molecule-Related/Downregulated by Oncogenes (*CDO*) and Brother of *CDO* (*BOC*) both bind Shh through Fn3 domains. Growth Arrest Specific 1 (*Gas1*) (Martinelli and Fan, 2007) and Hedgehog-Interacting Protein (*Hhip*) (Bosanac et al., 2009) interact with Shh, but not through Fn3 domains. Of these, expression of *CDO*, *BOC*, or *Gas1* increases Shh pathway activity, while *Hhip* negatively regulates the pathway (Beachy et al., 2010). SCUBE2, a secreted SCUBE protein family member, interacts with both Shh and Ptch and enhances Shh signaling (Tsai et al., 2009). The precise relationships between these receptor molecules and the Shh/Ptch interaction have yet to be described in detail.

p0140 The Gli transcription factors can also be regulated outside of the canonical Shh pathway. Borycki et al. demonstrated that Wnt1 and Wnt4 can induce *Gli2* expression and repress *Gli3* expression in a quail segmental plate mesoderm explant culture system (Borycki et al., 2000). Others have suggested that Gli1 protein may be regulated independently of Shh through the MAPK pathway (Seto et al., 2009). This would raise some interesting possibilities for activating Gli-regulated genes in the absence of Shh as well as for synergizing with Shh to superactivate the pathway.

p0145 While precise definitions of noncanonical pathways are lacking, results of multiple perturbations of the Shh pathway support the involvement of many of its components in multiple signaling paradigms. Indeed, it would be surprising if the enormous range of Shh effects as mitogen and morphogen in essentially every tissue could be reduced to one relatively simple pathway with three transcription factor effectors. Elaboration of these additional pathways for Shh signaling

and their roles in specific processes will be a rich source for potential targets of therapeutic molecules that are fine-tuned to specific effects that are perturbed in disease states.

Phenotypes of Shh pathway mutants

s0025

Hedgehog signaling is a fundamental pathway p0150 involved in many aspects of prenatal development. Varied roles have been described from a number of studies in model organisms using constitutive and targeted gene “knockouts” in mice and chick/quail chimeras (Table 1). The first report of a constitutive Shh knockout demonstrated cephalic neural tube defects as early as day 8.5 of gestation (E8.5) that becomes more severe a day later, resulting in a markedly hypomorphic central nervous system (Chiang et al., 1996). Mutant embryos die around the time of birth and exhibit defects in development of heart, lung, kidney, and foregut in addition to the forming CNS. Many of the same phenotypes have been observed in embryos lacking *Smo*, an intermediate member of the Shh pathway that functions to positively regulate pathway activation, and embryos lacking Dispatched A (*mDispA*), a factor that is essential for efficient Shh release from cells producing it. Embryos lacking *Smo* or *mDispA* have somewhat more severe phenotypes resembling *Shh* and Indian hedgehog (*Ihh*) double knockout. *mDispA* and *Smo* are required for *Ihh* as well as for Shh signaling.

Similar results are seen after exposure to an p0155 inhibitor of Shh signaling, cyclopamine, the effective agent in *Veratrum californicum* that induces cyclopia in fetuses of pregnant sheep. In addition to cyclopia, malformations of the nose and skull, notably the premaxilla, are also present (Binns et al., 1963). From these studies, it is clear that successful embryo development requires restriction of Shh activation to specific levels in both a temporal and spatial manner.

t0005 Table 1. Phenotypes caused by alterations and interruptions in *Shh* signaling that may relate to deficits in DS

Perturbation	Affected system	Phenotypes	Age	Notes	References
<i>Shh</i> ^{-/-}	Cardiac	Pharyngeal arch artery defects, ASD, VSD, Tetralogy of Fallot-like, persistent truncus arteriosus (PTA)	E10.5–15.5		Hildreth et al. (2009) and Smoak et al. (2005)
	Brain	Midline fusion of anterior lips of cephalic neural plate, incomplete separation of primitive optic vesicles ^a	E8.25		Abdelwahid et al. (2002)
		Loss of ventral structures, growth deficit in forebrain	E11.5		Chiang et al. (1996)
	Branchial arches	Reduced mandibular component ^a	E9.5		Abdelwahid et al. (2002)
	Craniofacial bones	Trace	E15.5		Chiang et al. (1996)
<i>Nkx2.5</i> ^{Cre/+} , <i>Shh</i> ^{fllox/-}	Cardiac	Pharyngeal arch artery defects, PTA, AVSD	E10.5	<i>Shh</i> ablated in <i>Nkx2.5</i> -expressing cells	Goddeeris et al. (2007)
<i>Mef2c-AHF-Cre</i> , <i>Smo</i> ^{fllox/-}	Cardiac	PTA, ASD, VSD, AVSD, rounded and short AV valves	E14.5–18.5	<i>Smo</i> ablated in anterior heart field	Goddeeris et al. (2008)
<i>Wnt1-Cre</i> , <i>Smo</i> ^{fllox/-}	Cardiac	PTA	E10.5	<i>Smo</i> ablated in neural crest	Goddeeris et al. (2007)
<i>Smo</i> ^{Gli1-CreERT2}	Cardiac	ASD and AVSD	E13.5	Floxed <i>Smo</i> allele under the control of inducible <i>Gli1:Cre</i>	Hoffmann et al. (2009)
<i>Shh</i> ^{Nkx2.1-Cre}	Cardiac	ASD	E13.5	<i>Shh</i> ablated in <i>Nkx2.1</i> -expressing cells	Hoffmann et al. (2009)
<i>Shh</i> ^{c/Shh} ⁿ , <i>Pax2-Cre</i>	Cerebellum	Absence of EGL, disorganized PL, fewer lobes	P5	<i>Shh</i> ablated in <i>Pax2</i> expressing precursors to Purkinje cells	Lewis et al. (2004)
<i>Shh</i> ^{c/Shh} ⁿ , <i>L7-Cre</i>	Cerebellum	Absence of EGL, disorganized PL, fewer lobes	P5	<i>Shh</i> ablated in precursors to Purkinje cells	Lewis et al. (2004)
<i>SuFu</i> ^{-loxP} , <i>Hoxb.7-Cre</i>	Cerebellum	Hypoplastic vermis, lack of foliation, disorganized cell layers	P21	<i>SuFu</i> ablated in precursors to CGNPs	Kim et al. (2011)
<i>5E1 Hybridoma</i>	Neural crest/branchial arches	Hypomorphic branchial arches, developmental delay	Stages 9–11 +24h	Hybridoma cells injected lateral mesenchyme of developing chick	Ahlgren and Bronner-Fraser (1999)
<i>Kif3a</i> ^{fl/fl} , <i>hGFAP-Cre</i>	Cerebellum	Atrophic cerebellum, fused folia, thicker PL	P25	<i>Kif3a</i> ablated in GFAP expressing CGNPS, Bergman glia, and Radial glia	Spassky et al. (2008)
<i>Ptc</i> ^{-/-}	Embryo	Failure of neural tube closure, lethality	E10.5	Constitutive <i>Ptc</i> knockout	Goodrich et al. (1997)
<i>Gli2</i> ^{fllox/fllox} , <i>En1-Cre</i>	Cerebellum	Reduced foliation, hypocellular EGL	P5/P8	<i>Gli2</i> is ablated by E9.0 in <i>En1</i> -expressing cells which give rise to the cerebellum	Blaess et al. (2006)

^aSimilar phenotypes were seen in *Smo* ^{-/-} and *mDispA* ^{-/-} knockout mice.

s0030 **Shh response deficit as a “common denominator” of DS phenotypes**

s0035 **Trisomy and Shh in cerebellar development**

p0160 The first direct demonstration of Shh response perturbation due to trisomy came from analysis of cerebellar development in the Ts65Dn “Down syndrome” mouse (Baxter et al., 2000). Mouse models play a critical role in the study of gene dosage mechanisms that produce the features of DS as reviewed in detail in this volume and elsewhere (Das and Reeves, 2011; de la Torre and Dierssen, 2012; Herault et al., 2012; Mobley, in this volume; Moore and Roper, 2007; O’Doherty et al., 2005). Ts65Dn mice, like people with trisomy 21, have a smaller cerebellum and show specific deficits of Purkinje cells and of the granule cell neurons that make up the internal granule layer (IGL) of the cerebellum. Further, the reduced density of GC in the IGL of Ts65Dn mice was shown to occur in people with DS, as well (Baxter et al., 2000).

p0165 The IGL is not present at birth in mice nor is it fully formed in newborn humans. Rather, granule cell precursors (GCPs) form the external germinal layer on the surface layer of the cerebellum. ~~It forms over~~ the first 3 weeks of life in mice (2–3 years in human beings); Purkinje cells produce Shh which stimulates GCPs to divide and migrate inward to form the IGL (Dahmane and Ruiz i Altaba, 1999; Wallace, 1999; Wechsler-Reya and Scott, 1999). The granule cell neuron deficit in Ts65Dn is already detectable from 1 week after birth (Roper et al., 2006b). On the day of birth, the number of GCPs in the external germinal layer is the same in Ts65Dn and euploid mice; however, the frequency of mitosis is significantly reduced in Ts65Dn. This reduced mitotic rate is a major contributor to the deficit in granule cell generation in trisomic mice (Roper et al., 2006b) and in DS (Guidi et al., 2011). Similarly, deleting a floxed Shh gene in late gestation by driving Cre expression with either the *Pax2* or *L7* promoters results in reduced cerebellar

volume, hypocellularity, and disorganization of GCPs in the EGL (Table 1; Lewis et al., 2004).

When GCPs were isolated from trisomic and euploid cerebella and cultured in the presence of increasing amounts of Shh, two important things were observed (Roper et al., 2006b). First, trisomic GCP responded less to the mitogenic effects of Shh at every concentration (Fig. 2a). Second, the trisomic cells did exhibit a dosage response, suggesting that stimulation of Shh signaling *in vivo* might overcome some of the mitogenic deficit in trisomic cells that was observed *in vitro*.




Fig. 2. SAG corrects Ts65Dn cerebellar dysmorphology. (a) Ts65Dn GCPs display an attenuated response to Shh treatment over a range of concentrations. (b) A single dose of SAG given on the day of birth restores GCP number at P6 to similar levels as Euploid littermates. (This figure is derived from Roper et al., 2006b.)

This was indeed the case. Trisomic mice that received a single dose of SAG on the day of birth had the same number of GCPs and of mitotic GCPs 4 week later, whereas vehicle-treated trisomic mice already showed a significant deficit of these cells (Fig. 2b).

s0040 **The *SHH* hypothesis for DS**

p0175 These results raise the question, “is the attenuated response to Shh in trisomic mice restricted to GCP, or do all Shh-responsive cells in a trisomic individual show a reduced reaction to Shh stimulation?” If the latter is the case, could stimulation of those developing populations at the appropriate stages of development represent a common approach to ameliorate diverse structural deficits in a wide range of cells and tissues that are affected to produce the DS phenotype? Based on the demonstration that trisomy results in a reduced response to the mitogenic effects of Shh in cerebellum, we consider here the possible effects of attenuated Shh response in three additional systems that are frequently or always affected during development in DS: craniofacial skeleton, heart, and the enteric nervous system. Observations of parallel effects of Shh disruption and of trisomy suggest that this mechanism may contribute to multiple DS phenotypes. Effects in development of the face and enteric nervous system further suggest that Shh signaling effects may impinge on neural crest cells (NCCs) which contribute to each of these structures.

s0045 **Craniofacial development**

p0180 The appearance of the DS face is very characteristic of this syndrome and is due substantially to hypoplasia of the midface skeleton. This is reflected in the Ts65Dn mouse and other models in an absolute correspondence between affected bones across the two species (Richtsmeier et al., 2000, 2002). In particular, the midface and mandible

are significantly smaller and dysmorphic due to trisomy. These bones arise from an embryonic precursor, Meckel’s cartilage, which is itself a product of the NCCs that contribute substantially to the first pharyngeal arch (PA1). To identify the earliest changes that lead to midface hypoplasia, we studied the formation of PA1 in Ts65Dn mice and their euploid littermates at embryonic day 9.5 (E9.5) in crosses with mice that express lacZ under control of the *Wnt1* promoter, marking NCCs (Roper et al., 2009).

Development of both trisomic and euploid p0185 embryos was highly variable at E9.5 with somite numbers ranging from 7 to 43, but no difference in developmental stage was observed between trisomic and euploid embryos (Roper et al., 2006a). When embryos at the 20–24 somite stage were considered, the trisomic PA1 was smaller, contained fewer neural crest-derived cells and these cells had a lower mitotic index than did their euploid counterparts. The number of migrating NCCs (lacZ + cells between the neural tube and PA1) was not significantly different at this stage; however, fewer migrating NCCs were present in slightly less mature, 17–19 somite embryos (Roper et al., 2009). Earlier experiments in both chick and mice show that Shh from endoderm of the ventral foregut is required to maintain migrating NCCs and to promote proliferation in PA1 (Ahlgren and Bronner-Fraser, 1999; Brito et al., 2006; Jeong et al., 2004). Note that if trisomic NCCs, like GCP, respond less to Shh than their euploid counterparts, some of these migrating cells might differentiate since they would “see” less Shh signal at the same concentration.

We then dissected the neural tubes from trisomic or euploid embryos and cultured them p0190 *ex vivo* to determine whether the delamination of NCCs from the tube is affected. Twenty-four hours after being placed in culture, Ts65Dn explants showed fewer cells migrating from the neural tube and those trisomic cells that did delaminate migrated for a shorter distance. Finally, we isolated cells from PA1 of euploid or trisomic embryos and cultured them to measure proliferation. Trisomic cells showed lower proliferation than did euploid.

However, addition of Shh to the cultures increased cell division, bringing the rate in trisomic cells to that seen in euploid cultures (Roper et al., 2006a).

p0195 Thus, the earliest trisomy-related deficits leading to midface skeletal hypoplasia arise from reduced delamination and migration of NCCs and from reduced proliferation of these cells in PA1, which provides the anlage for the cartilaginous model from which the mandible and mid-facial bones will form. The known effects of Shh as well as our observations are consistent with the hypothesis that an important contribution to this deficit is the reduced responsiveness to Shh in cranial neural crest from trisomic mice, with direct phenotypic consequences.

s0050 ***Trisomy and Shh in cardiovascular development***

p0200 Nearly half of all children born with DS have a congenital heart defect (CHD) (Ferencz et al., 1989). Atrioventricular septal defects (AVSDs) are the most common followed by ventricular septal defects (VSDs) and atrial septal defects (ASDs). Several mouse models of DS show similar patterns of CHD, indicating conservation of the effects of trisomic genes during mammalian heart development (Liu et al., 2011; O'Doherty et al., 2005; Williams et al., 2008). Nearly half of Tc1 transchromosomic mice, which carry a freely segregating copy of Hsa21, present with heart defects. VSDs are most common in trisomic mice, while AVSD and patent truncus arteriosus (PTA) are also observed. About 15% of newborn Ts65Dn mice have cardiac defects, including ASD, VSD, PTA, and various errors of branching of the pulmonary and outflow tracts. Mice that carry a duplication of the Hsa21 conserved synteny ~~region~~ on mouse chromosome 16 (Mmu16) show cardiovascular defects reminiscent of those seen in individuals with DS. They display ASD, VSD, and a tetralogy of Fallot-like phenotype (Li et al., 2007). Mice that are trisomic for all regions conserved with Hsa21 on mouse chromosomes 10, 16, and 17 have cardiovascular defects at a similar frequency (Yu et al., 2010).

Shh is secreted from cells in both the pulmonary endoderm, where it is required for proper atrial septation, and in the pharyngeal endoderm, where it is necessary for proper outflow tract septation (Goddeeris et al., 2008). Shh signaling marks cells within the second heart field (SHF) as progenitors of the atrial septum and outflow tract. Labeling of hedgehog-responsive cells early in heart development demonstrates that those cells migrate from the SHF and contribute to the primary atrial septum, dorsal mesenchymal protrusion (DMP), endocardial cushions, and pulmonary trunk (Hoffmann et al., 2009). The atrial septum, DMP, and endocardial cushions all combine to form the mesenchymal complex of the atrioventricular septum (Snarr et al., 2007). The appearance of this complex is necessary to complete AV septation and to anchor AV valves.

NCCs contribute to heart development by migrating into the outflow tract of the heart, contributing to septation and alignment. *Smo* is necessary for Shh pathway activation, and the loss of this gene in NCCs resulted in errors in septation and alignment of the aorta and pulmonary trunk, as well as defects in pharyngeal arch arteries (Goddeeris et al., 2008). A Shh response deficit could thus contribute to heart defects through direct effects in SHF, or because of an impaired response of trisomic neural crest. As noted, several steps in NCC delamination, migration, and proliferation require Shh signaling.

In support of this idea, several mouse models with impaired Shh signaling also display errors in septation (Table 1). A knockout of *Shh*^{-/-} in which exon 2 and its flanking introns are removed displays AVSD and other structural defects (Hildreth et al., 2009; Smoak et al., 2005). Similarly, when Shh signaling is blocked by cyclopamine at HH stage 14 chick embryos, they exhibit PTA, VSD, and pulmonary atresia secondary to reduced proliferation in the SHF (Dyer and Kirby, 2009). Similar outcomes occur when other components of the pathway are altered. Conditional knockouts of *Smo* and *Shh* result in AVSD and PTA in mouse embryos. Deletion of a floxed *Shh* allele in all

cells expressing either *Nkx2.5* or *Gli1* results in AVSD (Goddeeris et al., 2007; Hoffmann et al., 2009). Thus Shh signaling mutants present AVSDs, VSDs, and ASDs, structural defects that are common in DS (Ferencz et al., 1989).

p0220 Septal defects were attributed primarily to errors in the endocardial cushions for many years, but evidence has emerged recently that points to a critical role for DMP as a contributing factor, especially to AVSD and secundum ASD (Goddeeris et al., 2008; Hoffmann et al., 2009). In this light, it is relevant that Shh signaling is not required for endocardial cushion contributions to septation but is necessary for proper contributions to DMP from the SHF. When Shh signaling is disrupted in ~~DMP progenitors or the SHF~~, the DMP is hypoplastic or does not form and an AVSD results (Goddeeris et al., 2008; Hoffmann et al., 2009). Hypoplastic DMP has also been described in human fetuses with DS and in mice trisomic for all of *Mmu16* (Blom et al., 2003; Snarr et al., 2007; Webb et al., 1999). Thus there is an important role for Shh signaling in formation of the DMP, and for DMP involvement in AVSDs; DS is a major risk factor for AVSD (Ferencz et al., 1989). Overall, there are substantial similarities between heart phenotypes caused by trisomy and those seen in Shh signaling mutants. These results do not prove causation but they are consistent with the effects expected from reduced response to Shh signaling in the developing heart.

s0055 **Enteric nervous system**

p0225 The small and large intestines are innervated by vagal NCCs that migrate along the primitive gut from the rostral toward the caudal end in response to glial derived neurotrophic factor (GDNF) (Young et al., 2001). In humans, these enteric neuron precursors (ENPs) colonize the gut beginning week 7 of gestation, with the primitive enteric ganglia reaching the rectum in week 12 (Kenny et al., 2010). Failure of the ENPs to reach the caudal end of the colon results in a

condition known as aganglionic megacolon, or Hirschsprung's disease (HSCR) (Kenny et al., 2010). Though still rare, the incidence of HSCR in conjunction with DS is significantly increased over the rate in the population at large (Arnold et al., 2009). Mutations in the mouse *Ret* gene, a receptor tyrosine kinase that is activated by GDNF, cause NCCs colonizing the gut to migrate less efficiently and these mutants phenocopy HSCR (Asai et al., 2006). Human *RET* gene mutations contribute to susceptibility to the development of HSCR in people (Amiel et al., 2008; Angrist et al., 1995).

Shh is expressed by epithelial cells on the inner membrane of the gut and signals via BMP4 to inhibit differentiation of ENPs that are located in the central mesenchyme but are not close to the (outer) surface mesenchyme (Sukegawa et al., 2000). Inhibition of ENP differentiation could result in HSCR and, given the increased incidence in DS, it appears plausible that dosage-sensitive genes located on Hsa21 may contribute to the aganglionic phenotype. Decreased responsiveness to Shh could result in the expansion of the pro-differentiation environment to a point deeper in the gut mesenchyme than normal. Early differentiation of these ENPs could then deplete the migratory pool of cells before the entire length of colon has been colonized.

Hsa21 genes and Shh signaling

None of the genes encoding canonical Shh signaling pathway components are encoded on Hsa21. However, upregulation of *Ptch* (resulting in downregulation of the ~~SHH~~ pathway) has been reported in Ts65Dn mice for a specific, small group of stem cells in the subventricular zone (SVZ), the origin of granule cells in the dentate gyrus (Trazzi et al., 2011). In cultured neurospheres developed from the SVZ region, a C-terminal fragment of the APP protein, AICD, can contribute to the upregulation of *Ptch* transcription (Trazzi et al., 2011). Since the *APP* gene

is found on Hsa21 and thus is chronically upregulated in DS (and also in Ts65Dn mice), this provides a possible explanation for the attenuated mitogenic response to Shh by trisomic cells. At the phenotypic level, the number of cells in dentate gyrus is reduced by about 20% in Ts65Dn mice compared to euploid (Insausti et al., 1998; Lorenzi and Reeves, 2006). Drugs developed for AD that modulate APP cleavage to reduce C-terminal fragments might thus have an additional ameliorative benefit in DS.

p0240 Molecular pathway analysis has implicated
[Au7] several additional Hsa21 genes whose expression may impinge on Shh signaling directly or indirectly, especially on the regulation of *Gli1*,
[Au8] 2, and/or 3 (see review by Sturgeon et al., 2012). To date, however, there is no direct demonstration of a dosage-sensitive trisomic gene disrupting Shh signaling in the developing cerebellum, heart, or the cranial or vagal neural crest. Trisomic mouse models of DS provide a sensitized genetic background for dissection of these mechanisms.

s0065 Discussion

p0245 Trisomy for Hsa21 results in increased dosage for more than 300 genes, and numerous studies of gene expression in DS and in animal models suggest that most of these will be upregulated by ~50% whenever and wherever they are normally expressed. Viewed from this perspective, the challenge of finding “cures” based on the modulation of individual gene function is daunting. The availability of segmental trisomies in animal models that recreate the dosage imbalances seen in DS and the demonstration that this produces features analogous to those in DS (Reeves et al., 1995) have led to a productive phenotype-based approach to the development of therapies (Reeves and Garner, 2007; Reeves et al., 1995; Salehi et al., 2007)

p0250 The phenotype-based approach suggests the possibility that multiple effects of trisomy in different

tissues may result from perturbations in the same developmental pathways and regulatory processes, as we posit here for Shh. A deficit in response to the mitogenic effects of Shh has been demonstrated in trisomic cerebellar GCP. Trisomic NCC-derived cells in PA1 also appear to respond less to Shh than do their euploid counterparts. A similar response deficit in other trisomic cell types could affect development of the face, heart, enteric nervous system, and perhaps other tissues affected in DS. The cerebellar GCP response deficit to Shh is amenable to amelioration through the application of a small-molecule agonist of the Shh pathway (Chen et al., 2002; Roper et al., 2006b). Might a similar positive effect be possible in other cells and tissues that develop abnormally in DS if the Shh pathway could be stimulated to an appropriate degree at the appropriate time and place?

The Shh pathway is utilized in so many aspects p0255 of development that suggesting it as a therapeutic target seems highly improbable at first glance. Development is substantially disrupted in mice that are engineered to over- or underexpress Shh. In some case, the effects are concentration dependent, as when Shh acts as a morphogen to program cell response based on temporal and spatial gradients in anterior–posterior patterning of the limb (Harfe et al., 2004). Indeed, delivery of any molecule that stimulates or inhibits this pathway would likely need to be strictly limited in space and time to avoid deleterious side effects.

However, Shh should have no effect on cells p0260 that do not possess appropriate receptor and signal transduction pathways. We argue here that many if not all trisomic cells that are Shh responsive might show the attenuated response seen in cerebellar GCP. To the degree that this is the case, off-target effects would be reduced and could possibly have a beneficial effect.

The basic tenets of this model are testable in p0265 cell and mouse model systems. While it is clear that there is a substantial amount to learn about Shh signaling in all situations where it occurs, models of DS can play an important part in

understanding these pathways. If this single molecular mechanism does prove to be a “common denominator” of multiple trisomic phenotypes, there are attendant prospects that a single kind of pharmaceutical treatment might ameliorate multiple features of DS.

Acknowledgments

p0270 Critical support for this work was provided by the Down Syndrome Research and Treatment Foundation and Research Down Syndrome. This work was also supported by PHS awards 1R01 HD038384 from the National Institute of Child Health and Human Development and R01 HL083300 from the National Heart, Lung and Blood Institute. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Abbreviations

ACTB	Arizona Cognitive Test Battery
AD	Alzheimer disease
ASD	atrial septal defect
AVSD	atrioventricular septal defect
DMP	dorsal mesenchymal protrusion
DS	Down syndrome
EGL	external germinal layer of the cerebellum
ENP	enteric neuron precursors
GCP	granule cell precursor
HSCR	Hirschsprung's disease
IGL	internal granule layer
NCCs	neural crest cells
PA1	first pharyngeal arch
PTA	patent tricus arteriosus
SAG	Sonic agonist
SHF	second heart field
Shh	Sonic hedgehog
SVZ	subventricular zone
VSD	ventricular septal defect

References

- Abdelwahid, E., Pelliniemi, L. J., & Jokinen, E. (2002). Cell death and differentiation in the development of the endocardial cushion of the embryonic heart. *Microscopy Research and Technique*, 58, 395–403.
- Ahlgren, S. C., & Bronner-Fraser, M. (1999). Inhibition of sonic hedgehog signaling in vivo results in craniofacial neural crest cell death. *Current Biology*, 9, 1304–1314.
- Amiel, J., Sproat-Emison, E., Garcia-Barcelo, M., Lantieri, F., Burzynski, G., Borrego, S., et al. (2008). Hirschsprung disease, associated syndromes and genetics: A review. *Journal of Medical Genetics*, 45, 1–14.
- Angrist, M., Bolk, S., Thiel, B., Puffenberger, E. G., Hofstra, R. M., Buys, C. H., et al. (1995). Mutation analysis of the RET receptor tyrosine kinase in Hirschsprung disease. *Human Molecular Genetics*, 4, 821–830.
- Arnold, S., Pelet, A., Amiel, J., Borrego, S., Hofstra, R., Tam, P., et al. (2009). Interaction between a chromosome 10 RET enhancer and chromosome 21 in the Down syndrome-Hirschsprung disease association. *Human Mutation*, 30, 771–775.
- Asai, N., Fukuda, T., Wu, Z., Enomoto, A., Pachnis, V., Takahashi, M., et al. (2006). Targeted mutation of serine 697 in the Ret tyrosine kinase causes migration defect of enteric neural crest cells. *Development (Cambridge, England)*, 133, 4507–4516.
- Ayers, K. L., & Therond, P. P. (2010). Evaluating Smoothened as a G-protein-coupled receptor for Hedgehog signalling. *Trends in Cell Biology*, 20, 287–298.
- Barnes, E. A., Kong, M., Ollendorff, V., & Donoghue, D. J. (2001). Patched1 interacts with cyclin B1 to regulate cell cycle progression. *The EMBO Journal*, 20, 2214–2223.
- Barnfield, P. C., Zhang, X., Thanabalasingham, V., Yoshida, M., & Hui, C. C. (2005). Negative regulation of Gli1 and Gli2 activator function by Suppressor of fused through multiple mechanisms. *Differentiation*, 73, 397–405.
- Baxter, L. L., Moran, T. H., Richtsmeier, J. T., Troncoso, J., & Reeves, R. H. (2000). Discovery and genetic localization of Down syndrome cerebellar phenotypes using the Ts65Dn mouse. *Human Molecular Genetics*, 9, 195–202.
- Beachy, P. A., Hymowitz, S. G., Lazarus, R. A., Leahy, D. J., & Siebold, C. (2010). Interactions between Hedgehog proteins and their binding partners come into view. *Genes & Development*, 24, 2001–2012.
- Binns, W., James, L. F., Shupe, J. L., & Everett, G. (1963). A congenital cyclopi-type malformation in lambs induced by maternal ingestion of a range plant, *Veratrum californicum*. *American Journal of Veterinary Research*, 24, 1164–1175.
- Blaess, S., Corrales, J. D., & Joyner, A. L. (2006). Sonic hedgehog regulates Gli activator and repressor functions with spatial and temporal precision in the mid/hindbrain region. *Development (Cambridge, England)*, 133, 1799–1809.

- Blom, N. A., Ottenkamp, J., Wenink, A. G., & Gittenberger-de Groot, A. C. (2003). Deficiency of the vestibular spine in atrioventricular septal defects in human fetuses with down syndrome. *The American Journal of Cardiology*, 91, 180–184.
- Borycki, A., Brown, A. M., & Emerson, C. P. Jr. (2000). Shh and Wnt signaling pathways converge to control Gli gene activation in avian somites. *Development (Cambridge, England)*, 127, 2075–2087.
- Bosanac, I., Maun, H. R., Scales, S. J., Wen, X., Lingel, A., Bazan, J. F., et al. (2009). The structure of SHH in complex with HHIP reveals a recognition role for the Shh pseudo active site in signaling. *Nature Structural and Molecular Biology*, 16, 691–697.
- Braudeau, J., Delatour, B., Duchon, A., Lopes-Pereira, P., Dauphinot, L., de Chaumont, F., et al. (2011). Specific targeting of the GABA-A receptor [alpha]5 subtype by a selective inverse agonist restores cognitive deficits in Down syndrome mice. *Journal of Psychopharmacology*, 25, 1030–1042. [Au9]
- Brito, J. M., Teillet, M. A., & Le Douarin, N. M. (2006). An early role for sonic hedgehog from foregut endoderm in jaw development: Ensuring neural crest cell survival. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 11607–11612.
- Chen, J. K., Taipale, J., Young, K. E., Maiti, T., & Beachy, P. A. (2002). Small molecule modulation of Smoothened activity. *Proceedings of the National Academy of Sciences of the United States of America*, 99, 14071–14076.
- Chiang, C., Litingtung, Y., Lee, E., Young, K. E., Corden, J. L., Westphal, H., et al. (1996). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature*, 383, 407–413.
- Dahmane, N., & Ruiz i Altaba, A. (1999). Sonic hedgehog regulates the growth and patterning of the cerebellum. *Development (Cambridge, England)*, 126, 3089–3100.
- Das, I., & Reeves, R. H. (2011). The use of mouse models to understand and improve cognitive deficits in Down syndrome. *Disease Models & Mechanisms*, 4, 596–606.
- de la Torre, R., & Dierssen, M. (2012). Therapeutic approaches in the improvement of cognitive performance in Down syndrome: Past, present and future. *Progress in Brain Research*, 197, xxx–xxx. [TS1]
- Dyer, L. A., & Kirby, M. L. (2009). The role of secondary heart field in cardiac development. *Developmental Biology*, 336, 137–144.
- Edgin, J. O., Mason, G. M., Allman, M. J., Capone, G. T., Deleon, I., Maslen, C., et al. (2010). Development and validation of the Arizona Cognitive Test Battery for Down syndrome. *Journal of Neurodevelopmental Disorders*, 2, 149–164.
- Edgin, J. O., Mason, G. M., Spanò, G., Fernández, A., & Nadel, L. (2012). Human and mouse model cognitive phenotypes in Down syndrome: Implications for assessment. *Progress in Brain Research*, 197, xxx–xxx. [TS1]
- Ferencz, C., Neill, C. A., Boughman, J. A., Rubin, J. D., Brenner, J. I., & Perry, L. W. (1989). Congenital cardiovascular malformations associated with chromosome abnormalities: An epidemiologic study. *The Journal of Pediatrics*, 114, 79–86.
- Fernandez, F., Morishita, W., Zuniga, E., Nguyen, J., Blank, M., Malenka, R. C., et al. (2007). Pharmacotherapy for cognitive impairment in a mouse model of Down syndrome. *Nature Neuroscience*, 10, 411–413. [Au9]
- Fillat, C., & Altafaj, X. (2012). Gene therapy for Down syndrome. *Progress in Brain Research*, 197, xxx–xxx. [TS1]
- Goddeeris, M. M., Rho, S., Petiet, A., Davenport, C. L., Johnson, G. A., Meyers, E. N., et al. (2008). Intracardiac septation requires hedgehog-dependent cellular contributions from outside the heart. *Development (Cambridge, England)*, 135, 1887–1895.
- Goddeeris, M. M., Schwartz, R., Klingensmith, J., & Meyers, E. N. (2007). Independent requirements for Hedgehog signaling by both the anterior heart field and neural crest cells for outflow tract development. *Development (Cambridge, England)*, 134, 1593–1604.
- Goodrich, L. V., Milenković, L., Higgins, K. M., & Scott, M. P. (1997). Altered neural cell fates and medulloblastoma in mouse patched mutants. *Science (New York, NY)*, 277, 1109–1113.
- Guidi, S., Ciani, E., Bonasoni, P., Santini, D., & Bartesaghi, R. (2011). Widespread proliferation impairment and hypocellularity in the cerebellum of fetuses with Down syndrome. *Brain Pathology*, 21, 361–373.
- Hanson, J. E., Blank, M., Valenzuela, R. A., Garner, C. C., & Madison, D. V. (2007). The functional nature of synaptic circuitry is altered in area CA3 of the hippocampus in a mouse model of Down's syndrome. *The Journal of Physiology*, 579, 53–67.
- Harfe, B. D., Scherz, P. J., Nissim, S., Tian, H., McMahon, A. P., & Tabin, C. J. (2004). Evidence for an expansion-based temporal Shh gradient in specifying vertebrate digit identities. *Cell*, 118, 517–528.
- Haydar, T., & Reeves, R. H. (2012). Direct and indirect effects of trisomy on early brain development. *Trends in Neurosciences*, Submitted.
- Herault, Y., Duchon, A., Velot, E., Maréchal, D., & Brault, V. (2012). The in vivo Down syndrome genomic library in mouse. *Progress in Brain Research*, 197, xxx–xxx. [TS1]
- Hildreth, V., Webb, S., Chaudhry, B., Peat, J. D., Phillips, H. M., Brown, N., et al. (2009). Left cardiac isomerism in the Sonic hedgehog null mouse. *Journal of Anatomy*, 214, 894–904.
- Hoffmann, A. D., Peterson, M. A., Friedland-Little, J. M., Anderson, S. A., & Moskowitz, I. P. (2009). Sonic hedgehog is required in pulmonary endoderm for atrial septation. *Development (Cambridge, England)*, 136, 1761–1770.
- Humke, E. W., Dorn, K. V., Milenkovic, L., Scott, M. P., & Rohatgi, R. (2010). The output of Hedgehog signaling is

- controlled by the dynamic association between Suppressor of Fused and the Gli proteins. *Genes & Development*, 24, 670–682.
- Insausti, A. M., Megias, M., Crespo, D., Cruz-Orive, L. M., Dierssen, M., Vallina, I. F., et al. (1998). Hippocampal volume and neuronal number in Ts65Dn mice: A murine model of Down syndrome. *Neuroscience Letters*, 253, 175–178.
- Jenkins, D. (2009). Hedgehog signalling: Emerging evidence for non-canonical pathways. *Cellular Signalling*, 21, 1023–1034.
- Jeong, J., Mao, J., Tenzen, T., Kottmann, A. H., & McMahon, A. P. (2004). Hedgehog signaling in the neural crest cells regulates the patterning and growth of facial primordia. *Genes & Development*, 18, 937–951.
- Kenny, S. E., Tam, P. K., & Garcia-Barcelo, M. (2010). Hirschsprung's disease. *Seminars in Pediatric Surgery*, 19, 194–200.
- Kim, J. J., Gill, P. S., Rotin, L., van Eede, M., Henkelman, R. M., Hui, C.-c., et al. (2011). Suppressor of fused controls mid-hindbrain patterning and cerebellar morphogenesis via GLI3 repressor. *The Journal of Neuroscience*, 31, 1825–1836.
- Kleschevnikov, A. M., Belichenko, P. V., Villar, A. J., Epstein, C. J., Malenka, R. C., & Mobley, W. C. (2004). Hippocampal long-term potentiation suppressed by increased inhibition in the Ts65Dn mouse, a genetic model of Down syndrome. *The Journal of Neuroscience*, 24, 8153–8160.
- Kogerman, P., Grimm, T., Kogerman, L., Krause, D., Unden, A. B., Sandstedt, B., et al. (1999). Mammalian suppressor-of-fused modulates nuclear-cytoplasmic shuttling of Gli-1. *Nature Cell Biology*, 1, 312–319.
- Lewis, P. M., Gritli-Linde, A., Smeyne, R., Kottmann, A., & McMahon, A. P. (2004). Sonic hedgehog signaling is required for expansion of granule neuron precursors and patterning of the mouse cerebellum. *Developmental Biology*, 270, 393–410.
- Li, Z., Yu, T., Morishima, M., Pao, A., LaDuca, J., Conroy, J., et al. (2007). Duplication of the entire 22.9 Mb human chromosome 21 syntenic region on mouse chromosome 16 causes cardiovascular and gastrointestinal abnormalities. *Human Molecular Genetics*, 16, 1359–1366.
- Liu, C., Morishima, M., Yu, T., Matsui, S. I., Zhang, L., Fu, D., et al. (2011). Genetic analysis of Down syndrome-associated heart defects in mice. *Human Genetics*, 130, 623–632.
- Lorenzi, H. A., & Reeves, R. H. (2006). Hippocampal hypocellularity in the Ts65Dn mouse originates early in development. *Brain Research*, 1104, 153–159.
- Lott, I. T., & Dierssen, M. (2010). Cognitive deficits and associated neurological complications in individuals with Down's syndrome. *Lancet Neurology*, 9, 623–633.
- Mann, R. K., & Beachy, P. A. (2004). Novel lipid modifications of secreted protein signals. *Annual Review of Biochemistry*, 73, 891–923.
- Marigo, V., Davey, R. A., Zuo, Y., Cunningham, J. M., & Tabin, C. J. (1996). Biochemical evidence that patched is the Hedgehog receptor. *Nature*, 384, 176–179.
- Martinelli, D. C., & Fan, C. M. (2007). Gas1 extends the range of Hedgehog action by facilitating its signaling. *Genes & Development*, 21, 1231–1243.
- Moore, C. S., & Roper, R. J. (2007). The power of comparative and developmental studies for mouse models of Down syndrome. *Mammalian Genome*, 18, 431–443.
- O'Doherty, A., Ruf, S., Mulligan, C., Hildreth, V., Errington, M. L., Cooke, S., et al. (2005). An aneuploid mouse strain carrying human chromosome 21 with down syndrome phenotypes. *Science*, 309, 2033–2037.
- Pennington, B. F., Moon, J., Edgin, J., Stedron, J., & Nadel, L. (2003). The neuropsychology of Down syndrome: Evidence for hippocampal dysfunction. *Child Development*, 74, 75–93.
- Reeves, R. H., & Garner, C. C. (2007). A year of unprecedented progress in Down syndrome basic research. *Mental Retardation and Developmental Disabilities Research Reviews*, 13, 215–220.
- Reeves, R., Irving, N., Moran, T., Wohn, A., Kitt, C., Sisodia, S., et al. (1995). A mouse model for Down syndrome exhibits learning and behaviour deficits. *Nature Genetics*, 11, 177–183.
- Richtsmeier, J., Baxter, L., & Reeves, R. (2000). Parallels of craniofacial maldevelopment in Down Syndrome and Ts65Dn mice. *Developmental Dynamics*, 217, 137–145.
- Richtsmeier, J. T., Zumwalt, A., Carlson, E. J., Epstein, C. J., & Reeves, R. H. (2002). Craniofacial phenotypes in segmentally trisomic mouse models for Down syndrome. *American Journal of Medical Genetics*, 107, 317–324.
- Roper, R. J., Baxter, L. L., Saran, N. G., Klinedinst, D. K., Beachy, P. A., & Reeves, R. H. (2006). Defective cerebellar response to mitogenic Hedgehog signaling in Down [corrected] syndrome mice. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 1452–1456.
- Roper, R., St. John, H., Philip, J., Lawler, A., & Reeves, R. (2006). Perinatal loss of Ts65Dn mice, a model of Down syndrome. *Genetics*, 172, 437–443.
- Roper, R. J., VanHorn, J. F., Cain, C. C., & Reeves, R. H. (2009). A neural crest deficit in Down syndrome mice is associated with deficient mitotic response to Sonic hedgehog. *Mechanisms of Development*, 126, 212–219.
- Rueda, N., Florez, J., & Martinez-Cue, C. (2008). Chronic pentylentetrazole but not donepezil treatment rescues spatial cognition in Ts65Dn mice, a model for Down syndrome. *Neuroscience Letters*, 433, 22–27.
- Salehi, A., Delcroix, J. D., Belichenko, P. V., Zhan, K., Wu, C., Valletta, J. S., et al. (2006). Increased App expression in a mouse model of Down's syndrome disrupts NGF transport and causes cholinergic neuron degeneration. *Neuron*, 51, 29–42.

- Salehi, A., Faizi, M., Belichenko, P. V., & Mobley, W. C. (2007). Using mouse models to explore genotype-phenotype relationship in Down syndrome. *Mental Retardation and Developmental Disabilities Research Reviews*, 13, 207–214.
- Salehi, A., Faizi, M., Colas, D., Valletta, J., Laguna, J., Takimoto-Kimura, R., et al. (2009). Restoration of norepinephrine-modulated contextual memory in a mouse model of Down syndrome. *Science Translational Medicine*, 1, 7ra17.
- Seto, M., Ohta, M., Asaoka, Y., Ikenoue, T., Tada, M., Miyabayashi, K., et al. (2009). Regulation of the hedgehog signaling by the mitogen-activated protein kinase cascade in gastric cancer. *Molecular Carcinogenesis*, 48, 703–712.
- Smock, I. W., Byrd, N. A., Abu-Issa, R., Goddeeris, M. M., Anderson, R., Morris, J., et al. (2005). Sonic hedgehog is required for cardiac outflow tract and neural crest cell development. *Developmental Biology*, 283, 357–372.
- Snarr, B. S., O'Neal, J. L., Chintalapudi, M. R., Wrigg, E. E., Phelps, A. L., Kubalak, S. W., et al. (2007). Isl1 expression at the venous pole identifies a novel role for the second heart field in cardiac development. *Circulation Research*, 101, 971–974.
- Spassky, N., Han, Y.-G., Aguilar, A., Strehl, L., Besse, L., Laclef, C., et al. (2008). Primary cilia are required for cerebellar development and Shh-dependent expansion of progenitor pool. *Developmental Biology*, 317, 246–259.
- Stone, D. M., Hynes, M., Armanini, M., Swanson, T. A., Gu, Q., Johnson, R. L., et al. (1996). The tumour-suppressor gene patched encodes a candidate receptor for Sonic hedgehog. *Nature*, 384, 129–134.
- Sturgeon, X., Le, T., Ahmed, M. M., & Gardiner, K. J. (2012). Pathways to cognitive deficits in Down syndrome. *Progress in Brain Research*, 197, xxx–xxx.
- Sukegawa, A., Narita, T., Kameda, T., Saitoh, K., Nohno, T., Iba, H., et al. (2000). The concentric structure of the developing gut is regulated by Sonic hedgehog derived from endodermal epithelium. *Development (Cambridge, England)*, 127, 1971–1980.
- Thibert, C., Teillet, M. A., Lapointe, F., Mazelin, L., Le Douarin, N. M., & Mehlen, P. (2003). Inhibition of neuroepithelial patched-induced apoptosis by sonic hedgehog. *Science*, 301, 843–846.
- Trazzi, S., Mitrugno, V. M., Valli, E., Fuchs, C., Rizzi, S., Guidi, S., et al. (2011). APP-dependent up-regulation of Pthch1 underlies proliferation impairment of neural precursors in Down syndrome. *Human Molecular Genetics*, 20, 1560–1573.
- Tsai, M. T., Cheng, C. J., Lin, Y. C., Chen, C. C., Wu, A. R., Wu, M. T., et al. (2009). Isolation and characterization of a secreted, cell-surface glycoprotein SCUBE2 from humans. *The Biochemical Journal*, 422, 119–128.
- Wallace, V. A. (1999). Purkinje-cell-derived Sonic hedgehog regulates granule neuron precursor cell proliferation in the developing mouse cerebellum. *Current Biology*, 9, 445–448.
- Wang, B., Fallon, J. F., & Beachy, P. A. (2000). Hedgehog-regulated processing of Gli3 produces an anterior/posterior repressor gradient in the developing vertebrate limb. *Cell*, 100, 423–434.
- Webb, S., Anderson, R. H., Lamers, W. H., & Brown, N. A. (1999). Mechanisms of deficient cardiac septation in the mouse with trisomy 16. *Circulation Research*, 84, 897–905.
- Wechsler-Reya, R. J., & Scott, M. P. (1999). Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron*, 22, 103–114.
- Williams, A. D., Mjaatvedt, C. H., & Moore, C. S. (2008). Characterization of the cardiac phenotype in neonatal Ts65Dn mice. *Developmental Dynamics*, 237, 426–435.
- Young, H. M., Hearn, C. J., Farlie, P. G., Canty, A. J., Thomas, P. Q., & Newgreen, D. F. (2001). GDNF is a chemoattractant for enteric neural cells. *Developmental Biology*, 229, 503–516.
- Yu, T., Li, Z., Jia, Z., Clapcote, S. J., Liu, C., Li, S., et al. (2010). A mouse model of Down syndrome trisomic for all human chromosome 21 syntenic regions. *Human Molecular Genetics*, 19, 2780–2791.
- Yue, S., Chen, Y., & Cheng, S. Y. (2009). Hedgehog signaling promotes the degradation of tumor suppressor Sufu through the ubiquitin-proteasome pathway. *Oncogene*, 28, 492–499.
- Zeng, X., Goetz, J. A., Suber, L. M., Scott, W. J., Jr., Schreiner, C. M., & Robbins, D. J. (2001). A freely diffusible form of Sonic hedgehog mediates long-range signalling. *Nature*, 411, 716–720.
- Zhang, C., Williams, E. H., Guo, Y., Lum, L., & Beachy, P. A. (2004). Extensive phosphorylation of Smoothened in Hedgehog pathway activation. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 17900–17907.

B978-0-444-54299-1.00011-X, 00011

Author Query Form

Book Series: Progress in Brain Research, 197
Chapter 11



Dear Author,

During the preparation of your manuscript for typesetting some questions have arisen. These are listed below. Please check your typeset proof carefully and mark any corrections in the margin of the proof or compile them as a separate list. This form should then be returned with your marked proof/list of corrections to Elsevier Science.

Disk use

In some instances we may be unable to process the electronic file of your article and/or artwork. In that case we have, for efficiency reasons, proceeded by using the hard copy of your manuscript. If this is the case the reasons are indicated below:

- ☐ Disk damaged ☐ Incompatible file format ☐ LaTeX file for non-LaTeX journal
☐ Virus infected ☐ Discrepancies between electronic file and (peer-reviewed, therefore definitive) hard copy.
☐ Other:

We have proceeded as follows:

- ☐ Manuscript scanned ☐ Manuscript keyed in ☐ Artwork scanned
☐ Files only partly used (parts processed differently:.....)

Bibliography

If discrepancies were noted between the literature list and the text references, the following may apply:

- ☐ The references listed below were noted in the text but appear to be missing from your literature list. Please complete the list or remove the references from the text.
☐ Uncited references: This section comprises references which occur in the reference list but not in the body of the text. Please position each reference in the text or, alternatively, delete it. Any reference not dealt with will be retained in this section.

Query Refs.	Details Required	Author's response
Au1	Please provide telephone and fax number to the corresponding author.	
Au2	Please check whether "patent truncus arteriosus" can be changed to "persistent truncus arteriosus" in the abbreviation list.	
Au3	Please check the section levels.	
Au4	Please include the details for "Mobley (in this volume)."	
Au5	Please note that "Nadel (in this volume)" has been changed to "Edgin et al. (2012)" as per TOC.	
Au6	Please check the clarity of the sentence "When Shh signaling..."	
Au7	As per scientific conventions, genes have to be styled in italics and proteins in roman. Please mark in the proofs for the same.	
Au8	Please note that "Gardiner (in this volume)" has been changed to "Sturgeon et al. (2012)" as per TOC.	
Au9	Please check the inserted volume and page range for this reference.	
TS1	We will update the page numbers in the revises/final stage.	